


Search for ultra-high energy photons with the Telescope Array surface detector

Mikhail Kuznetsov, Oleg Kalashev and Grigory Rubtsov
for the Telescope Array collaboration

ICRC-2019
Madison, US
July 29, 2019




Supported by Russian Science Foundation

- ▶ Telescope Array experiment
- ▶ Technique of photon search — multivariate analysis
- ▶ Blind search for point sources of photons

based on Abbasi et al., arXiv:1904.00300

- ▶ Target search for photons from dwarf galaxies


Telescope Array experiment: surface detector





- ▶ 507 SD's, 3 m² each
- ▶ 680 km² area
- ▶ 11 years of operation

Largest UHECR statistics in the Northern Hemisphere

p -induced EAS



γ -induced EAS



Photon-induced showers:

- ▶ arrive younger
- ▶ contain less muons
- ▶ multiple SD observables affected: **front curvature**, **Area-over-peak**, $\chi^2/d.o.f.$, etc.

Data and Monte-Carlo sets

- ▶ Data collected by TA surface detector for the 9 years:

2008-05-11 — 2017-05-10

- ▶ p and γ Monte-Carlo sets with CORSIKA and dethinning

Stokes et al, Astropart.Phys.35:759,2012

Cuts for both data and MC:

- ▶ 7 or more detectors triggered
- ▶ core distance to array boundary is larger than 1200m
- ▶ $\chi^2/\text{d.o.f.} < 5$
- ▶ $\theta < 60^\circ$
- ▶ $E_\gamma > 10^{18}$ eV (E_γ is estimated with photon Monte-Carlo)
- ▶ remove events coincident with lightnings (lightning events mimics γ -induced events)

52362 events after cuts

Photon search: list of relevant SD observables





1. Linsley front curvature parameter, a ;
2. Area-over-peak (AoP) of the signal at 1200 m;
Pierre Auger Collaboration, Phys.Rev.Lett. 100 (2008) 211101
3. AoP LDF slope parameter;
4. Number of detectors hit;
5. N. of detectors excluded from the fit of the shower front;
6. $\chi^2/d.o.f.$;
7. $S_b = \sum S_i \times r^b$ parameter for $b = 3$ and $b = 4.5$;
Ros, Supanitsky, Medina-Tanco et al. Astropart.Phys. 47 (2013) 10
8. The sum of signals of all detectors of the event;
9. Asymmetry of signal at upper and lower layers of detectors;
10. Total n. of peaks within all FADC traces;
11. N. of peaks for the detector with the largest signal;
12. N. of peaks present in the upper layer and not in lower;
13. N. of peaks present in the lower layer and not in upper;

Machine learning for multivariate analysis.

- ▶ The Boosted Decision Trees (BDT) technique is used to build p - γ classifier based on multiple observables.
Telescope Array, Astropart. Phys. 110, 8 (2019); PRD 99, 022002 (2019)
- ▶ root::TMVA is used as a stable implementation.
PoS ACAT 040 (2007), arXiv:physics/0703039
- ▶ BDT is trained with Monte-Carlo sets: γ (signal) and p (background)*
- ▶ BDT classifier is used to convert the set of observables of each event to a number $\xi \in [-1 : 1]$
- ▶ ξ is available for one-dimensional analysis.

* MC set is split into 3 equal parts: (I) for training the classifier, (II) for ξ -cut optimization, (III) for exposure estimate.

Distribution of MVA estimator ξ for data and MC



data proton MC gamma MC

Search for point sources of photons: motivation

The way to improve the photon search sensitivity:

Hadron background is highly isotropic



Assume that photons are emitted by point source



In angular vicinity of the source the **photon/hadron ratio** would be larger than in full TA field of view



Easier to separate photons from hadrons!

Bonus!

Specific photon source hypotheses could be tested by search in certain directions stacked

- Dwarf spheroidal galaxies (heavy DM decay hypothesis)

Search for point sources of photons: motivation

The way to improve the photon search sensitivity:

Hadron background is highly isotropic



Assume that photons are emitted by point source



In angular vicinity of the source the **photon/hadron ratio** would be larger than in full TA field of view



Easier to separate photons from hadrons!

Bonus!

Specific photon source hypotheses could be tested by search in certain directions stacked

- Dwarf spheroidal galaxies (heavy DM decay hypothesis)

Features of γ point sources search: blind search

- ▶ Independent search for γ in each skymap direction
- ▶ The angular size of the each search region is equal to the γ **angular resolution**:


$E_\gamma \geq$, eV	$10^{18.0}$	$10^{18.5}$	$10^{19.0}$	$10^{19.5}$	$10^{20.0}$
ang.res.	3.00°	2.92°	2.64°	2.21°	2.06°

- ▶ The skymap is pixelized into 12288 directions with HEALpix (7868 in TA field of view)

Optimisation of MVA-cut for γ flux upper-limit:

- ▶ Assume the flux consists of protons only (null hypothesis):
 $F_{\text{total}} = F_p$
- ▶ Optimize the ξ -cut separately for the best upper-limit in each direction using MC p and MC γ
- ▶ E^{-2} γ -spectrum is assumed

Results: point-source photon flux upper-limits




$E_\gamma \geq, \text{eV}$	$\langle F_\gamma \rangle \leq, \text{km}^{-2}\text{yr}^{-1}$
$10^{18.0}$	0.094
$10^{18.5}$	0.029
$10^{19.0}$	0.010
$10^{19.5}$	0.0071
$10^{20.0}$	0.0058

Pierre Auger: $\langle F_\gamma \rangle \leq 0.035 \text{ km}^{-2}\text{yr}^{-1}$ (1° ang.res., $10^{17.3} \leq E \leq 10^{18.5} \text{ eV}$)

A. Aab et al. ApJ 789, 160 (2014)

Results: point-source photon flux upper-limits

95% C.L.




The results for all points are available in table form with paper

Abbasi et al., arXiv:1904.00300

It can be used to constrain models of UHECR sources.


Results: photon excesses significance



$E_\gamma \geq, \text{eV}$	max. γ signif. (pre-trial)
$10^{18.0}$	2.72σ
$10^{18.5}$	2.71σ
$10^{19.0}$	2.89σ
$10^{19.5}$	2.76σ
$10^{20.0}$	3.43σ

The excesses are insignificant, given the large number of trials

Results: photon excess significance



Though not significant enough, the largest excess is 3.43σ for $E_\gamma \geq 10^{20.0}$ eV located at $\{\alpha = 155.3^\circ, \delta = 60.4^\circ\}$

Target search for photons from dwarf galaxies

Probe for the possible decay of heavy dark matter (HDM)

- ▶ HDM decay produce significant amount of photons in any model
M. Kachelriess et al., PRD 98, 083016 (2018)
- ▶ DM is abundant in dwarf galaxies (Galactic Center is outside the TA field of view)
- ▶ Target source set: 21 dwarf galaxies — satellites of Milky Way
V. Bonnivard et al., MNRAS 453 (2015), 849
- ▶ Search for γ in stacked skymap pixels of dwarf galaxies (pixel size = γ **ang.res.**)

Results

No evidence for photon signal ($N_{\gamma}^{\text{cand.}} = 0$ at all energies)

E_{γ}, eV	$10^{18.0}$	$10^{18.5}$	$10^{19.0}$	$10^{19.5}$	$10^{20.0}$
$F_{\text{UL}}^{\gamma}, \text{km}^{-2}\text{yr}^{-1}$	0.15	0.057	0.014	0.0076	0.0052

These results can be used to constrain HDM models

Target search for photons from dwarf galaxies

Probe for the possible decay of heavy dark matter (HDM)

- ▶ HDM decay produce significant amount of photons in any model
M. Kachelriess et al., PRD 98, 083016 (2018)
- ▶ DM is abundant in dwarf galaxies (Galactic Center is outside the TA field of view)
- ▶ Target source set: 21 dwarf galaxies — satellites of Milky Way
V. Bonnivard et al., MNRAS 453 (2015), 849
- ▶ Search for γ in stacked skymap pixels of dwarf galaxies (pixel size = γ **ang.res.**)

Results


No evidence for photon signal ($N_{\gamma}^{\text{cand.}} = 0$ at all energies)

E_{γ}, eV	$10^{18.0}$	$10^{18.5}$	$10^{19.0}$	$10^{19.5}$	$10^{20.0}$
$F_{UL}^{\gamma}, \text{km}^{-2}\text{yr}^{-1}$	0.15	0.057	0.014	0.0076	0.0052


These results can be used to constrain HDM models

Plans

Improve γ -search sensitivity with recently developing Neural Net




Waveform encoder:



see poster #172 on Tuesday

Increase γ -search exposure with recently deploying TAx4 SD detector



► **Blind photon search**

- The directional upper-limits for photons with $E > 10^{18}$ eV are set
- The directional upper-limits for $E > 10^{18.5}$ eV are set for the first time
- No significant evidence for photon signal was found in blind search
- The results can be used to constrain UHECR sources models

► **Target photon search**

- The photons from staked dwarf galaxies were searched — no candidates were found
- The flux upper-limits for dwarf galaxies can be used to constrain heavy dark matter models

Thank you!

► **Blind photon search**

- The directional upper-limits for photons with $E > 10^{18}$ eV are set
- The directional upper-limits for $E > 10^{18.5}$ eV are set for the first time
- No significant evidence for photon signal was found in blind search
- The results can be used to constrain UHECR sources models

► **Target photon search**

- The photons from staked dwarf galaxies were searched — no candidates were found
- The flux upper-limits for dwarf galaxies can be used to constrain heavy dark matter models

Thank you!

Backup slides

Monte-Carlo: photon angular resolution

The geometry reconstruction of events is crucial for point source search.

There is a bias in zenith angle θ reconstruction.

We correct all data and MC events for the mean value of this bias:

$$\langle \theta_{\text{rec.}} - \theta_{\text{true}} \rangle$$

Angular reconstruction for photon primaries (used in this search)

E_γ , eV	$\langle \theta_{\text{rec.}} - \theta_{\text{true}} \rangle$	ang. resolution
$10^{18.0}$	-2.25°	3.00°
$10^{18.5}$	-2.24°	2.92°
$10^{19.0}$	-2.16°	2.64°
$10^{19.5}$	-2.06°	2.21°
$10^{20.0}$	-1.72°	2.06°

Features of γ point sources search: blind search

- ▶ Independent search for γ in each skymap direction
- ▶ The angular size of the search region is equal to the γ **angular resolution**
- ▶ The skymap is pixelized into 12288 directions with HEALpix (7868 in TA field of view)

Optimisation of MVA-cut for γ flux upper-limit:

- ▶ Assume the flux consists of protons only (null hypothesis):
 $F_{\text{total}} = F_p$
- ▶ Optimize the ξ -cut separately in each constant declination band
 $\delta_j \pm \text{ang.res.}$ using MC p and MC γ

- ▶ The optimization is for the minimum upper-limit:

$$F_{\gamma}^{\text{UL}} = \frac{\mu_{\text{Poisson}}^{\text{UL}}(N_p^{\xi})}{A_{\gamma}^{\text{eff}}}, \quad \text{where } N_p^{\xi} = \{\text{number of MC } p \text{ events passing the } \xi\text{-cut}\};$$

$\mu_{\text{Poisson}}^{\text{UL}} = \{\text{upper-limit of the Poisson mean}\}; A_{\gamma}^{\text{eff}} = \{\text{Effective exposure of the experiment to } \gamma \text{ for the given values of event quality cuts and } \xi\text{-cut}\}$